



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROFESSOR ROEMER, of Marburg, has been called to Greifswald to conduct the hygienic institute as the successor of Professor Loeffler.

DISCUSSION AND CORRESPONDENCE

A NEW TYPE OF BACTERIAL DISEASE

By this title I mean a disease in which the bacterial growth first develops conspicuously as a thick layer on the surface of the plant, and only later penetrates into its interior.

Ráthay's disease of orchard grass (*Dactylis glomerata*) described by him in 1899 may be taken as the type of this kind of disease. In 1913 I had opportunity to verify Ráthay's statements¹ on material sent to me from Denmark by Professor Kølpin Ravn, and to make pure cultures and further studies of the organism which in honor of Ráthay, may be known as *Aplanobacter ráthayi* n. sp., with the characters assigned to it by Ráthay, and in addition the following:

Nitrates are not reduced; gelatin is finally liquefied, but liquefaction is visible only after some weeks and progresses very slowly; it does not grow in Cohn's solution; growth starts off slowly in milk, but is prolonged with formation of a copious chrome yellow precipitate and a wide bright yellow rim; litmus milk is first slowly blued, but becomes purplish after some weeks; it grows so slowly on agar that poured plates which appear to be sterile may eventually give small yellow colonies. Nearly all of Ráthay's statements have been found to be correct. This note is here published because of delay in the issue of a longer account.

ERWIN F. SMITH

THE MANUS OF TRACHODONT DINOSAURS

IN a recent article in *The Ottawa Naturalist*,¹ Mr. Lawrence M. Lambe has described "The Manus in a Specimen of *Trachodon* from the Edmonton of Alberta," illustrated by three figures. According to Mr. Lambe's interpretation of the Ottawa skeleton the phalangeal formula is as follows:

¹ *Sitz. Ber. Wiener Akad.*, 1 Abt., Bd. CVIII., p. 597.

² Vol. XXVII., pp. 21-25, 1913.

Digit II. with three phalanges, the third bearing a hoof.

Digit III. with three phalanges, the third bearing a hoof.

Digit IV. with two phalanges, the second bearing a hoof.

Digit V. with two phalanges, the second bearing a hoof.

Whereas in a specimen that I have described the formula is

Digit II. with three phalanges, the third bearing a hoof.

Digit III. with three phalanges, the third bearing a hoof.

Digit IV. with three phalanges, the third a vestigial bone without hoof.

Digit V. with three phalanges, the third a vestigial bone without hoof.

The writer published a description of the manus of *Trachodon annectens*,² based on the first reported specimen in which all of the phalanges are present. In this specimen the full number of phalanges are not only present but each digit is articulated either in the right or the left hand and all are encased in a thin layer of matrix in which the skin impression is preserved.

In this uncrushed specimen the long slender metacarpals of digits II., III., and IV. are closely appressed as represented in the figure accompanying the above article, a position verified by structure and by position in three other uncrushed specimens in the American Museum, one in the National Museum, and a sixth in the collection of the Calgary Natural History Society.

In no specimen of the genus *Trachodon* known to me have more than two hoof bones been found in the manus—those of digits II. and III. The terminal phalanges of digits IV. and V. are, when uncrushed, rounded bony nodules, very much reduced and were not covered by a hoof or nail.

If Mr. Lambe's interpretation is correct we have a remarkable specific variation in this genus in which a later species, described by me, has developed an additional phalanx on each

² *Bull. Am. Mus. Nat. Hist.*, Vol. XXXI., Art. X., pp. 105-107, 1912.